

SHRIKRISHNA MAHA VIDYALAYA, GUNJOTI



Project Report on

**Existence of the freshwater fish faunal diversity in
Osmanabad district (MS), India**

Submitted By

Jawan Sneha Dhondiba, Seat No: NAF664921
Mhetre Akash Gangadhar, Seat No: NAF664942
Potdar Sairaj Balaji, Seat No: NAF664955
Rathod Omkar Gulab, Seat No: NAF664957
Rathod Balika Ashok, Seat No: NAF664956

Guided By

Dr. D.M.Pathan
Asso. Prof. Dept. of Zoology

Submitted to


Shrikrishna shikshan sanstha's

SHRIKRISHNA MAHAVIDYALAYA GUNJOTI


DEPARTMENT OF ZOOLOGY

CERTIFICATE


This is to certify that, Mr. /Miss Jawan S. D.,Mhetre A.G.,Potdar S.B.,rathod O.G.,Rathod B.A. Class B.Sc. VI Semester, Zoology has satisfactorily completed the Project on **Existence of the freshwater fish faunal diversity in Osmanabad district (MS), India** as per instructed by Dr. B. A. M. University, Aurangabad during the academic year 2021-22.



Guide



External Examiner



Head

HEAD

Department of Zoology
S. K. M. Gunjoti Tr. Omergo

Existence of the freshwater fish faunal diversity in Osmanabad district (MS), India

The present study deals with the existence of the fish faunal diversity in riverine and reservoir in Osmanabad district (Mah), India. During the investigation study period 2015-16 it was observed that the number of 26 species of fish fauna belonging to 12 families and 6 orders were recorded. The Cypriniformes represented 11 species of the fishes, followed by Siluriformes - 6 species, Perciformes - 5 species, Ostcoglossiformes - 2 species, Anguilliformes - 1 species and Synbranchiformes - 1 species were observed. The highest number of 11 species was recorded in the order of Cypriniformes. The predominant order of fish fauna in this district are Cypriniformes, Siluriformes and Perciformes. All above fish fauna species recorded were found to be widely distributed in the riverine and reservoirs. Detailed results are summarized in the present paper

INTRODUCTION : Biodiversity is an important factor for the stability of an ecosystem. In the Asian region the knowledge of the freshwater fish faunal biodiversity and its conservation aspects are relatively less documented as it is still in exploration and discovery phase (Nguyen and DeSilva, 2006). Similarly, Indian fish fauna remains in need of in depth systematic study as many species are still to be described or to be discovered and the available information is from a few well studied location only (Bhat, 2003., Goyal and Arora, 2009, Le`ve`que *et al.*, 2008; Molur and Walkar, 1998). India is one of the mega biodiversity countries in the world. There are rich in freshwater ecosystems (Kar *et al.*, 2006). There are 450 families of freshwater fishes globally, out of which 40 families are represented from India (Jayaram, 2010, Keshave *et al.*, 2013). India possesses maximum number of fresh-water fishes comprising 225 species (DeSilva, 2007; Karmarkar, and Das, 2005). The freshwater resources are currently experiencing an alarming rate of decline in fish diversity with 17 species critically endangered, 69 species under endangered and 81 species under vulnerable status in the East Himalayas and Western Ghats (Allen *et al.*, 2010; Molur *et al.*, 2011; Shinde *et al.*, 2009).

Dam and reservoirs in India, which are constantly increasing in number, play an important role not only in electric and water supply but also in providing a source of fish to be local community, for food, research, sustainable aquaculture and maintenance of fish diversity (Yusoff and Ambak, 1999). Most of the rivers and reservoirs are now dominated by exotic fish like grass carps, silver carp and other predatory fishes, which tolerate high pollution and static water levels. Over exploitation and habitat degradation as an example have depleted the stocks and reduced the replacement rate in the population (Khan *et al.*, 1996).

Previous study indicates that most of the work is related to fish fauna available from riverine and reservoir ecosystem. Very little attention is information about the freshwater fish diversity except the work (Kharat *et al.*, 2012; Shinde *et al.*, 2009). In the present investigation an attempt has been made to highlight the fish diversity to formulate future strategies for development and fish conservation & also helps in species selection for aquaculture in this region

MATERIAL AND METHODS

Osmanabad is one of the district of Marathwada regions of the state of Maharashtra in India. It is situated in the southern part of the state. It lies on the Deccan plateau, about 600 m above sea level. It lies between north latitudes 17° 35' and 18° 40' and longitude 75° 16' and 76° 40'. The Osmanabad district (fig-1) has a geographical area of 7512 sq.km the district forms part of Godavari basin and Manjra sub basin. Manjra, Sina, Terna, Bori, Benitura, Banganga are the main rivers flowing through the district.

The rainy season starts from mid-June and continues till the end of September. The climate is humid in October and November and dry and cool from mid-November to January. From February to June the climate is dry and becomes increasingly hot. During summer the temperature of Osmanabad district is low compared to other districts of Marathwada region. The average annual rainfall in the district is 730 mm. Temperature max 42.10c and min 80c. The freshwater fishes were collected with the help of local fisherman by fishing craft, gears and various types of nets are used (fig-2). Collected fishes were properly preserved in 10% formalin in laboratory. All the specimen fishes were identified performed standard literature. The Integrated Taxonomic Information System (ITIS) standard report fish base (<http://fishbase.org>) and other reference books are used as fish identified up to species level.

RESULTS AND DISCUSSION

In the present investigation total number of 26 species occurs in freshwater fish fauna belonging to 6 orders. The analysis of data indicated that the order Cypriniformes was dominant with 11 fish species followed by order Siluriformes 6 fish species and Perciformes 5 fish species. The order Osteoglossiformes two fish species, Anguilliformes and Synbranchiformes each with one species have been recorded are shown in Table 1 and Fig 3.

Studied Ichthyofish diversity and conservation aspect in a lake and rivers ecosystem in India's inland water resources are diversified as they are plentiful (Khan *et al.*, 1996; Kharat *et al.*, 2012). Reservoir contributed the single largest inland fishery resources both in terms of size and production potential (Kamble and Medkhede, 2013). Fish species were the important indicator of ecological health. The abundance and health of fish showed the health of water bodies (Hamzah, 2007).

The present work is concluded that the existence of freshwater fish fauna diversity assumes top most priority under changing habitat degradation and biological characters of species serve the baseline information for further studies on resource conservation and maintenance. The finding

of this study is expected to benefit the planning and management toward sustainable fishery and conservations programs of riverine and reservoir. The total number of species recorded during this study period has shown a good indicator of rich diversity in riverine and reservoir of Osmanabad district in Maharashtra.

Conflicts of interest: The authors stated that no conflicts of interest.

Table 1: The freshwater fish fauna diversity existence in various river and reservoirs of Osmanabad district. Sr. No	Order	Family	Scientific name of the fish species
1	Cypriniformes	Cyprinidae	<i>Catla catla</i> (Ham and Jhingran)
			<i>Cirrhinus mrigala</i> (Hamilton)
			<i>Cyprinus carpio</i> (Linnaeus)
			<i>Labeo rohita</i> (Ham and Buch)
			<i>Labeo calbasu</i> (Ham and Buch)
			<i>Labeo fambriatus</i> (Ham)
			<i>Puntius ticto</i> (Ham)
			<i>Puntius vittatus</i> (Day)
			<i>Rasbora daniconius</i> (Ham and Buch)
			<i>Ctenopgaryngodon idella</i>
			Nemacheilinae
2	Siluriformes	Bagridae	<i>Nemacheilus anguilla</i>
			<i>Mystus seenghala</i> (Sykes)
			<i>Mystus cavasius</i>
			<i>Mystus vittatus</i>
			Siluridae
			<i>Wallago attu</i> (Schneider)
			<i>Heteropneustes fossilis</i> (bloch)
			Heteropneustidae
			<i>Clarias batrachus</i> (Linnaeus)
			Claridae
3	Perciformes	Channidae	<i>Channa marulius</i> (Ham)
			<i>Channa punctatus</i> (Bloch)
			<i>Channa striatus</i> (Bloch)
			Anabantidae
			<i>Anabas testudineus</i>
			Cichlidae
			<i>Oreochromis mossambicus</i> (Peters)
4	Anguilliformes	Anguillidae	<i>Anguillia</i>

5		Notopteridae	<i>bengalensis</i> (Gray) <i>Notopterus</i> <i>notopterus</i> (Pallas)
	Osteoglossiformes		
	<i>Notopterus chitala</i> (Ham)		
6	Synbranchiformes	Mastacembelidae	<i>Mastacembelus</i> <i>armatus</i> (Lac)

REFERENCES

- Allen DJ, Molur S and Daniel BA (2010) The Status and Distribution of Freshwater Biodiversity in the Eastern Himalaya. 1-88. Published by IUCN, Cambridge, UK and Gland, Switzerland and Zoo Outreach organization, Coimbatore, India
- Bhat (2003) Diversity and composition of freshwater fishes in streams of Central Western Ghats, India. *Environmental Biology of Fishes*, 68: 25-38.
- De Silva Sena, Abery NW and Nguyen TTT (2007) Endemic freshwater finfish of Asia: Distribution and conservation status. *Diversity and Distribution*. 13:172-184.
- Ghatge SS and Shelke ST (2013) Inventory of endemic freshwater fish fauna of Maharashtra state: India. *Rec Zool. Survey. India*: 113 (part-3): 79-92.
- Goyal AK and Arora S (2009) India's Fourth National Report to the Convention on Biological Diversity, Ministry of Environment and Forests, *Govt. of India*. 1-143.
- Hamzah N (2007) Assessment on water quality and biodiversity within Sungai Batu Pahat. *Master of thesis, University Technology, Malaysia*, 124.
- Jayaram KC (2010) The fresh water fishes of Indian region. 2nd edition, *Narendra Publication*, New Delhi.
- Kamble AT and Medkhede LM (2013) Study of fish fauna and productivity of Loni reservoir, Tq. Kinwat (Maharashtra). *International Journal of Biomedical and Advance Research*. 4 (3): 155-159.
- Kar D, Nagarathna AV, Ramchandra TV and Dey SC (2006) Fish diversity and conservation aspects in an aquatic ecosystem in North Eastern India. *Zoo's print*, 21(7): 2308-2315.
- Karmarkar AK and Das A (2005) Endemic Freshwater Fishes of India. *Rec. Zool. Surv, India. Occ. Paper No.* 230:1-125.
- Keshave JV, Ananthan PS and Landge L (2013) Fish diversity and productivity of Isapur Reservoir, Maharashtra state. *International Journal of Biomedical and Advance Research*, 4(12): 865-867.
- Khan MS, Lee, Patrick KY, Cramphorn J and Zakaria Ismail Mohd (1996) Freshwater fishes of the Pahang River Basin, Malaysia. *Wetland International J. Asia Pacific Publication*. 112.
- Kharat SS, Dahanukar N, Raut R and Mahabaleshwarkar M (2003) Long-term changes in freshwater fish species composition in Northern Western Ghats, Pune district. *Current Science*, 84(6): 816-820.
- Kharat SS, Paingankar M and Dahanukar N (2012) Freshwater fish fauna of Krishna River at Wai, Northern Western Ghats of India. *Journal of Threatened Taxa*. 4(6): 2569-2577.



Fig. 2: Capturing the freshwater fish of the local fisherman in Osmanabad district (Mah).